

**IPEA**

Instituto de Planejamento Econômico e Social

PLAN

Instituto de Planejamento

TEXTO PARA DISCUSSÃO INTERNA

Nº 101

"Testing for First Order Serial  
Correlation in Temporally  
Aggregated Regression Models"

Pedro L. Valls Pereira

Novembro de 1986



TEXTO PARA DISCUSSÃO INTERNA  
Nº 101

"Testing for First Order Serial  
Correlation in Temporally  
Aggregated Regression Models"

Pedro L. Valls Pereira

Novembro de 1985

30.11.1985

17-12-85

Tiragem: 50 exemplares

Trabalho Concluído em: Novembro de 1986

Instituto de Pesquisas do IPEA  
Instituto de Planejamento Econômico e Social  
Avenida Presidente Antônio Carlos, 51 - 130/170 andares  
20.020 Rio de Janeiro RJ

Este trabalho é da inteira e exclusiva responsabilidade de seu autor. As opiniões nele emitidas não exprimem, necessariamente, o ponto de vista da Secretaria de Planejamento da Presidência da República.

SECRETARIA DE P. H.	
INSTITUTO DE PLANEJAMENTO ECONÔMICO E SOCIAL	
SEIOR DE I	TOMBO
	8693-2
F. N. 1029	
DATA 29, 04, 87	

# TESTING FOR FIRST ORDER SERIAL CORRELATION IN TEMPORALLY AGGREGATED REGRESSION MODELS

Pedro L. Valls Pereira<sup>1</sup>  
INPES - IPEA

## ABSTRACT

This paper shows that the LM statistic for testing first order serial correlation in regression models can be computed using the Kalman Filter.

It is shown that when there are missing observations, the LM statistic for this test is equivalent to the test statistic derived by Robinson (1985) using the likelihood conditional on the observation times.

The Kalman Filter approach is preferable because the test statistic for first order serial correlation in temporally aggregated regression models can be obtained as an extension of the previous case.

**KEYWORDS:** First Order Serial Correlation; LM test; Regression Models; Missing Observations; Temporal Aggregation; Kalman Filter.

---

<sup>1</sup>This paper was written during a visit to LSE. Research support was provided by the British Council and Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq). I am grateful to the members of the DEMEIC workshop for helpful comments and specially to Prof. Peter Robinson.

## 1 INTRODUCTION

The most common test against autocorrelated errors in regression models is the bounds test of Durbin & Watson (1950, 1951, 1971). Many authors - Breusch (1978) and Godfrey (1978) among others - observed that the Durbin's h statistic ( see Durbin (1970)) can be derived from the general theory of the LM test procedure.

When observations are missing in regression models, a modified Durbin - Watson statistic was proposed by Savin & White (1978) and Dufour & Dagenais (1985).

In a recent paper Robinson (1985) applied the LM principle and derived two test statistics, where one is obtained using an unconditional form of the likelihood and the other using the likelihood conditional on the observation times. Asymptotic distributions of the test statistics are established and analytic comparison of efficiencies are made.

The motivation for using the LM approach is that it produces an asymptotically locally most powerful test and the  $\chi^2$  asymptotic approximation to its distribution is robust to departures from normality.

This paper follows Robinson's approach and a LM statistic for testing serial correlation in regression models with missing observations is obtained. This test statistic is obtained as a by - product of the Kalman Filter recursions. It is shown that a reparametrization of the parameter space is useful in obtaining this test statistic.

This approach is used because the extension to temporally aggregated regression models can be done very easily.

The structure of this paper is as follows. In section 2, the LM statistic for testing serial correlation in regression models with missing observations is obtained.

Section 3 extends the results to temporally aggregated regression models.

Concluding remarks are made in Section 4.

## 2 THE LM STATISTIC FOR TESTING FIRST ORDER SERIAL CORRELATION IN REGRESSION MODELS WITH MISSING OBSERVATIONS

Consider the regression model with autocorrelated errors, i. e.:

$$y(t) = x'(t)\beta + u(t) \quad t = 1, \dots, T \quad (2.1)$$

$$u(t) = \rho u(t-1) + \epsilon(t) \quad (2.2)$$

where  $x(t)$  is a  $k \times 1$  vector of nonstochastic regressors, independent of  $u(t)$ ,  $\beta$  is a  $k \times 1$  vector of unknown parameters and  $\epsilon(t)$  is i.i.d.  $N(0, \sigma^2)$ .

It is assumed that  $y(t)$  is observed each  $m$  time periods (sometimes called "skip sampling"), i. e.  $y_\tau = y(m\tau)$   $\tau = 1, \dots, \lfloor \frac{T}{m} \rfloor$  is the observed endogenous variable and  $x(t)$  is observed for each time period.

The proposition below proves that the regression model for the observed data has parameters  $\beta$ ,  $\rho_*$  and  $\sigma_*^2 = (1 + \rho^2 + \dots + \rho^{2(m-1)})\sigma^2$ . Therefore the log likelihood function for this regression model will be a function of the new parameter set  $\Psi = (\rho_*, \beta', \sigma_*^2)$  and the LM statistic for testing  $H_0 : \rho = 0$  vs  $H_a : \rho \neq 0$  will be obtained using this reparametrization, because testing  $\rho = 0$  is equivalent to test  $\rho_* = 0$ .

**Proposition I.**

Consider the regression model with autocorrelated errors given by (2.1-2). If the endogenous variable  $y(t)$  is observed only for  $t \equiv 0 \pmod{m}$ , the observed model is given by:

$$y_\tau = x_\tau' \beta + u_\tau \quad \tau = 1, \dots, \lfloor \frac{T}{m} \rfloor \quad (2.3)$$

$$u_\tau = \rho_* u_{\tau-1} + \epsilon_\tau^* \quad (2.4)$$

and  $\epsilon_\tau^*$  is i. i. d.  $N(0, \sigma_*^2)$  where  $\sigma_*^2 = (1 + \rho^2 + \dots + \rho^{2(m-1)})\sigma^2$  and  $\rho_* = \rho^m$ .

**Proof:** see Appendix I

In order to obtain the log likelihood function for the model (2.3-4) the Kalman Filter recursions will be used.

The state space representation of the model (2.3-4) is given by the measurement equation, i. e.:

$$y_\tau = (1, 0) \alpha_\tau \quad \tau = 1, \dots, \lfloor \frac{T}{m} \rfloor \quad (2.5)$$

$$\Leftrightarrow y_\tau = Z_\tau \alpha_\tau$$

and the transition equation, i. e.:

$$\alpha_\tau = \begin{bmatrix} 0 & \rho_* \\ 0 & \rho_* \end{bmatrix} \alpha_{\tau-1} + \begin{bmatrix} x_\tau' \beta \\ 0 \end{bmatrix} + \begin{bmatrix} 1 \\ 1 \end{bmatrix} \epsilon_\tau^* \quad (2.6)$$

$$\Leftrightarrow \alpha_\tau = \Phi \alpha_{\tau-1} + c_\tau + R \epsilon_\tau^*$$

where  $\text{Var}(\epsilon_\tau^*) = \sigma_*^2 Q$ .

The log likelihood function can be obtained using the prediction error decomposition (see Harvey (1981)) and is given by:

$$l(\Psi) = \ln L(\rho_*, \beta', \sigma_*^2) = -\frac{T}{2m} \ln 2\pi - \frac{T}{2m} \ln \sigma_*^2$$

$$- \frac{1}{2} \sum_{\tau=1}^{\lfloor \frac{T}{m} \rfloor} \ln f_\tau - \frac{1}{2\sigma_*^2} \sum_{\tau=1}^{\lfloor \frac{T}{m} \rfloor} \frac{\nu_\tau^2}{f_\tau} \quad (2.7)$$

where  $\nu_\tau$  is the prediction error and  $\sigma_*^2 f_\tau$  is the variance of the prediction error and they are obtained by the Kalman Filter recursions (see Harvey (1981) pages 107 - 111). These recursions are given by two sets of equations: the prediction equations and the updating equations. In order to start these recursions, the mean,  $a_0$ , and the variance - covariance matrix,  $\sigma_*^2 P_0$ , of the state space vector at time  $t = 0$  are needed. The log likelihood (2.7) is obtained when these starting values are given by:

$$a_0 = 0$$

$$P_0 = \begin{bmatrix} \kappa & 0 \\ 0 & \frac{1}{1-\rho_*^2} \end{bmatrix}$$

where  $\kappa$  is a large number, e. g.  $\kappa = 10^6$ , which is equivalent to use a diffuse prior for the initial conditions (see Ansley & Kohn (1985)).

It is easy to see that

$$\nu_1 = y_1 - x_1' \beta \quad (2.8)$$

$$\nu_\tau = (y_\tau - x_\tau' \beta) - \rho_*(y_{\tau-1} - x_{\tau-1}' \beta) \quad \tau = 2, \dots, \left[ \frac{T}{m} \right] \quad (2.9)$$

and

$$f_1 = \frac{1}{1 - \rho_*^2} \quad (2.10)$$

$$f_\tau = 1 \quad \tau = 2, \dots, \left[ \frac{T}{m} \right] \quad (2.11)$$

In order to derive the LM statistic for testing for first order autocorrelation, the first derivatives of the log likelihood function and the information matrix are needed. Pagan (1978) or Engle & Watson (1981) have shown that the information matrix can be computed using only first derivatives of the likelihood function. These derivatives are computed using auxiliary recursions (see Appendix II).

By Breusch & Pagan (1978), the LM statistic for testing for first order autocorrelation is given by:

$$LM = \tilde{D}_1 \left( \tilde{I}_{\rho_* \rho_*} - \tilde{I}_{\rho_* \beta} \tilde{I}_{\beta \beta}^{-1} \tilde{I}_{\beta \rho_*} \right)^{-1} \tilde{D}_1 \quad (2.12)$$

where

$$\tilde{D} = \begin{bmatrix} \frac{\partial l}{\partial \rho_*} \\ \frac{\partial l}{\partial \beta} \\ \frac{\partial l}{\partial \sigma_*^2} \end{bmatrix} (\tilde{\Psi}) = \begin{bmatrix} \tilde{D}_1 \\ 0 \\ 0 \end{bmatrix}$$

and

$$\tilde{I} = \begin{bmatrix} I_{\rho_* \rho_*} & I_{\rho_* \beta} & I_{\rho_* \sigma_*^2} \\ I_{\beta \rho_*} & I_{\beta \beta} & I_{\beta \sigma_*^2} \\ I_{\sigma_*^2 \rho_*} & I_{\sigma_*^2 \beta} & I_{\sigma_*^2 \sigma_*^2} \end{bmatrix} (\tilde{\Psi})$$

is the Fisher information matrix evaluated at  $\tilde{\Psi} = \Psi_{H_0} = (0, \tilde{\beta}, \tilde{\sigma}_*^2)$  and

$$\begin{aligned} \tilde{\beta} &= [(X^*)' (X^*)]^{-1} (X^*)' Y^* \\ X^* &= (x(m), x(2m), \dots, x(T)) \\ Y^* &= (y(m), y(2m), \dots, y(T)) \\ \tilde{\sigma}_*^2 &= \frac{1}{T} \sum_{\tau=1}^{\left[ \frac{T}{m} \right]} (y_\tau - x_\tau' \tilde{\beta})^2 = \frac{1}{T} \sum_{\tau=1}^{\left[ \frac{T}{m} \right]} \tilde{u}_\tau^2 \end{aligned}$$

From Appendix III it follows that:

$$LM = \frac{T}{m} (r_1^*)^2 \quad (2.13)$$

where

$$r_1^* = \frac{\sum_{\tau=1}^{\lfloor \frac{T}{m} \rfloor} \tilde{u}_\tau \tilde{u}_{\tau-1}}{\sum_{\tau=1}^{\lfloor \frac{T}{m} \rfloor} \tilde{u}_\tau} \quad (2.14)$$

which is exactly the  $s_1$  statistic derived by Robinson (1985).

### 3 THE LM STATISTICS FOR TESTING FIRST ORDER SERIAL CORRELATION IN TEMPORALLY AGGREGATED REGRESSION MODELS

Now it is assumed that the endogenous variable  $y(t)$  in (2.1) is a flow variable and instead of observing  $y(t)$  every time period, is  $y_\tau$  which is observed, i. e.:

$$y_\tau = \sum_{j=0}^{m-1} y(m\tau - j) \quad \tau = 1, \dots, \left\lfloor \frac{T}{m} \right\rfloor \quad (3.1)$$

Given that model (2.1-2) is the disaggregated model, i. e. for times  $t = 1, \dots, t$ , Proposition II below derives the aggregated model, i. e. for  $\tau = 1, \dots, \left\lfloor \frac{T}{m} \right\rfloor$  and the endogenous variable is given by (3.1).

#### Proposition II.

Given model (2.1-2) and assume that  $y_\tau$  is given by (3.1). Then the aggregated model is given by:

$$y_\tau = z_\tau' \beta + u_\tau \quad \tau = 1, \dots, \left\lfloor \frac{T}{m} \right\rfloor \quad (3.2)$$

$$u_\tau = \rho_* u_{\tau-1} + \epsilon_\tau^* + \theta_* \epsilon_{\tau-1}^* \quad (3.3)$$

where

$$z_\tau' = \sum_{j=0}^{m-1} x'(m\tau - j)$$

$$\rho_* = \rho^m$$

$$\epsilon_\tau^* \sim N(0, \sigma_*^2)$$

and  $\theta_*$  and  $\sigma_*^2$  are given by the solutions of the following equations:

$$(1 + \theta_*^2) \sigma_*^2 = [1 + (1 + \rho)^2 + (1 + \rho + \rho^2)^2 + \dots + (1 + \rho + \dots + \rho^{m-1})^2 + \rho^2(1 + \rho + \dots + \rho^{m-2})^2 + \rho^4(1 + \rho + \dots + \rho^{m-3})^2 + \dots + \rho^{2(m-1)}] \sigma^2 \quad (3.4)$$

$$\theta_* \sigma_*^2 = [\rho(1 + \rho + \dots + \rho^{m-2}) + (1 + \rho)\rho^2(1 + \rho + \dots + \rho^{m-3}) + \dots + (1 + \rho + \dots + \rho^{m-3})\rho^{m-2}(1 + \rho) + (1 + \rho + \dots + \rho^{m-2})\rho^{m-1}] \sigma^2 \quad (3.5)$$

$$|\theta_*| < 1 \quad (3.6)$$



**Proof:** : see Appendix IV

The LM statistic for testing first order serial correlation will be derived using the Kalman Filter recursions. In order to use these recursions the state space representation of model (3.2-3) is needed.

This representation is given by the measurement equation, i. e.:

$$y_\tau = (0, 0, 1) \alpha_\tau \quad \tau = 1, \dots, \left[ \frac{T}{m} \right] \quad (3.7)$$

$$\Leftrightarrow y_\tau = Z_\tau \alpha_\tau$$

and the transition equation, i. e.:

$$\alpha_\tau = \begin{bmatrix} \rho_* & 1 & 0 \\ 0 & 0 & 0 \\ \rho_* & 1 & 0 \end{bmatrix} \alpha_{\tau-1} + \begin{bmatrix} 0 \\ 0 \\ z_\tau' \beta \end{bmatrix} + \begin{bmatrix} 1 \\ \theta_* \\ 1 \end{bmatrix} \epsilon_\tau^* \quad (3.8)$$

$$\Leftrightarrow \alpha_\tau = \Phi \alpha_{\tau-1} + c_\tau + R \epsilon_\tau^*$$

where  $\text{Var}(\epsilon_\tau^*) = \sigma_*^2 Q$ .

The Kalman Filter recursions are started by

$$a_0 = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

$$P_0 = \begin{bmatrix} \frac{1+2\rho_*\theta_*+\theta_*^2}{1-\rho_*^2} & \theta_* & 0 \\ \theta_* & \theta_*^2 & 0 \\ 0 & 0 & \kappa \end{bmatrix}$$

and the log likelihood function is given by (2.7).

It is easy to see that:

$$\nu_1 = y_1 - z_1' \beta \quad (3.9)$$

$$\nu_\tau = (y_\tau - z_\tau' \beta) - \rho_*(y_{\tau-1} - z_{\tau-1}' \beta) - \theta_* f_{\tau-1}^{-1} \nu_{\tau-1} \quad \tau = 2, \dots, \left[ \frac{T}{m} \right] \quad (3.10)$$

and

$$f_1 = \frac{1 + 2\rho_*\theta_* + \theta_*^2}{1 - \rho_*^2} \quad (3.11)$$

$$f_\tau = 1 + \theta_*^2 (1 - f_{\tau-1}^{-1}) \quad \tau = 2, \dots, \left[ \frac{T}{m} \right] \quad (3.12)$$

It is easy to see that under  $H_0$  ( and under some conditions on the  $x(t)$ ), i. e.  $\rho = 0 \Leftrightarrow \rho_* = 0$ :

(i) the OLS estimate of  $\beta$ , i. e.  $\tilde{\beta}$  is given by:

$$\tilde{\beta} = (Z'Z)^{-1} Z'Y^* \quad (3.13)$$

- where  $Z = (z_1, \dots, z_{\lfloor \frac{T}{m} \rfloor})$ ,  $z_\tau = \sum_{j=0}^{m-1} x(m\tau - j)$ ;
- (ii) the prediction error is given  $\nu_\tau = y_\tau - z_\tau' \tilde{\beta} \quad \forall \tau$ ;
  - (iii) the variance of the prediction error is given by  $f_\tau = 1 \quad \forall \tau$ ;
  - (iv) and  $\tilde{\sigma}_*^2 = m\tilde{\sigma}^2 = \frac{1}{m} \sum_{\tau=1}^{\lfloor \frac{T}{m} \rfloor} (y_\tau - z_\tau' \tilde{\beta})^2 = \frac{1}{m} \sum_{\tau=1}^{\lfloor \frac{T}{m} \rfloor} \tilde{u}_\tau^2$ .
- From Appendix V, it follows that:

$$LM = \frac{T}{m} (r_1^*)^2 \quad (3.14)$$

where

$$r_1^* = \frac{\sum_{\tau=1}^{\lfloor \frac{T}{m} \rfloor} \tilde{u}_\tau \tilde{u}_{\tau-1}}{\sum_{\tau=1}^{\lfloor \frac{T}{m} \rfloor} \tilde{u}_\tau} \quad (3.15)$$

#### 4 CONCLUDING REMARKS

It was shown that the LM statistic for testing first order serial correlation can be derived as a by-product of the Kalman Filter recursions.

In order to use the Kalman Filter recursions to derive the LM statistic a reparametrization was necessary.

It was shown that the LM statistic for testing first order serial correlation in regression models with missing observations, is equivalent to the  $s_1$  statistic derived by Robinson (1985). This statistic is obtained from the likelihood conditional on the observation times. The LM statistic can be interpreted as  $\lfloor \frac{T}{m} \rfloor$  times  $R^2$ , the coefficient of determination of the regression of  $\tilde{u}_\tau$  in  $\tilde{u}_{\tau-1}$ , where  $\tilde{u}_\tau = y_\tau - z_\tau'$ , i. e. the residuals, under  $H_0$ , of model (2.1) for every  $m$  time period. An extension for regression models where the endogenous variable is observed at every time period for  $t = 1, \dots, p_0 T$  ( $1 \leq p_0 < 1$ ) and for  $t = p_0 T + 1, \dots, T$  is observed at every  $m$  time period, is given in Pereira (1985).

Using the Kalman Filter recursions, it was possible to extend this result to temporally aggregated regression models. As before the LM statistic can be interpreted as  $\lfloor \frac{T}{m} \rfloor$  times  $R^2$  coefficient of determination of the regression of  $\tilde{u}_\tau$  in  $\tilde{u}_{\tau-1}$ , where  $\tilde{u}_\tau = y_\tau - z_\tau'$  and  $z_\tau = \sum_{j=0}^{m-1} x(m\tau - j)$ , i. e. the residuals under  $H_0$  of model (3.2-3).

#### APPENDIX I

(2.1-2) can be rewritten as follows:

$$(1 - \rho L)y(t) = (1 - \rho L)x'(t)\beta + \epsilon(t) \quad (A.I.1)$$

Multiplying both sides of (A.I.1) by  $(1 + \rho L + \dots + \rho^{m-1} L^{m-1})$ , it follows:

$$(1 - \rho^m L^m)y(t) = (1 - \rho^m L^m)x'(t)\beta + (1 + \rho L + \dots + \rho^{m-1} L^{m-1})\epsilon(t) \quad (A.I.2)$$

Now define

$$\begin{aligned} B &= L^m \\ \rho_* &= \rho^m \\ \epsilon_\tau^* &= (1 + \rho L + \dots + \rho^{m-1} L^{m-1}) \epsilon(t) \quad \tau = 1, \dots, \left[ \frac{T}{m} \right] \end{aligned}$$

then (A.I.2) can be rewritten as follows:

$$(1 - \rho_* B) y_\tau = (1 - \rho_* B) x_\tau' \beta + \epsilon_\tau^* \quad (\text{A.I.3})$$

⇕

$$y_\tau = x_\tau' \beta + u_\tau \quad (\text{A.I.4})$$

$$u_\tau = \rho_* u_{\tau-1} + \epsilon_\tau \quad (\text{A.I.5})$$

Using the results of Tiao(1972) or Pereira(1986) it follows that  $\epsilon_\tau^* \sim NID(0, \sigma_*^2)$  where  $\sigma_*^2 = (1 + \rho^2 + \dots + \rho^{2(m-1)}) \sigma^2$ .

## APPENDIX II

Taking derivatives of (2.3) with respect to the parameters, it follows that:

$$\begin{aligned} \frac{\partial l}{\partial \psi_i} &= \frac{1}{2\sigma_*^2} \sum_{\tau=1}^{\left[ \frac{T}{m} \right]} \frac{\nu_\tau^2}{f_\tau} \frac{\partial f_\tau}{\partial \psi_i} - \frac{1}{2\sigma_*^2} \sum_{\tau=1}^{\left[ \frac{T}{m} \right]} \frac{\nu_\tau}{f_\tau} \frac{\partial \nu_\tau}{\partial \psi_i} \\ &\quad - \frac{1}{2} \sum_{\tau=1}^{\left[ \frac{T}{m} \right]} \frac{1}{f_\tau} \frac{\partial f_\tau}{\partial \psi_i} \quad i = 1, \dots, k+1 \end{aligned} \quad (\text{A.II.1})$$

$$\frac{\partial l}{\partial \sigma_*^2} = \frac{T}{2m\sigma_*^2} + \frac{1}{2\sigma_*^4} \sum_{\tau=1}^{\left[ \frac{T}{m} \right]} \frac{\nu_\tau^2}{f_\tau} \quad (\text{A.II.2})$$

and the information matrix is given by:

$$I_{\psi_i \psi_j} = \frac{1}{\sigma_*^2} \sum_{\tau=1}^{\lfloor \frac{T}{m} \rfloor} \frac{1}{f_\tau} \begin{bmatrix} \partial \nu_\tau \\ \partial \psi_i \end{bmatrix} \begin{bmatrix} \partial \nu_\tau \\ \partial \psi_j \end{bmatrix} - \frac{1}{2} \sum_{\tau=1}^{\lfloor \frac{T}{m} \rfloor} \frac{1}{f_\tau^2} \frac{\partial f_\tau}{\partial \psi_i} \frac{\partial f_\tau}{\partial \psi_j} \quad i, j = 1, \dots, k+1 \quad (\text{A.II.3})$$

$$I_{\sigma_*^2 \psi_i} = \frac{1}{\sigma_*^4} \sum_{\tau=1}^{\lfloor \frac{T}{m} \rfloor} \frac{\nu_\tau}{f_\tau} \frac{\partial \nu_\tau}{\partial \psi_i} - \frac{1}{2\sigma_*^4} \sum_{\tau=1}^{\lfloor \frac{T}{m} \rfloor} \frac{\nu_\tau^2}{f_\tau} \frac{\partial f_\tau}{\partial \psi_i} \quad i = 1, \dots, k+1 \quad (\text{A.II.4})$$

$$I_{\sigma_*^2 \sigma_*^2} = \frac{T}{2m\sigma_*^4} \quad (\text{A.II.5})$$

In order to obtain (A.II.1-4), it is needed auxiliary recursions for the derivatives. They are:

#### Prediction Equations

$$\frac{\partial a_{\tau/\tau-1}}{\partial \psi_i} = \frac{\partial \Phi}{\partial \psi_i} a_{\tau-1} + \Phi \frac{\partial a_{\tau-1}}{\partial \psi_i} + \frac{\partial c_\tau}{\partial \psi_i} \quad (\text{A.II.6})$$

$$\begin{aligned} \frac{\partial P_{\tau/\tau-1}}{\partial \psi_i} &= \frac{\partial \Phi}{\partial \psi_i} P_{\tau-1} \Phi' + \Phi \frac{\partial P_{\tau-1}}{\partial \psi_i} \Phi' + \Phi P_{\tau-1} \left( \frac{\partial \Phi}{\partial \psi_i} \right)' \\ &+ \frac{\partial R}{\partial \psi_i} Q R' + R \frac{\partial Q}{\partial \psi_i} R' + R Q \left( \frac{\partial R}{\partial \psi_i} \right)' \quad i = 1, \dots, k+1 \end{aligned} \quad (\text{A.II.7})$$

#### Prediction Error

$$\frac{\partial f_\tau}{\partial \psi_i} = Z_\tau' \frac{\partial P_{\tau/\tau-1}}{\partial \psi_i} Z_\tau \quad i = 1, \dots, k+1 \quad (\text{A.II.8})$$

#### Variance of the Prediction Error

$$\frac{\partial \nu_\tau}{\partial \psi_i} = -Z_\tau' \frac{\partial a_{\tau/\tau-1}}{\partial \psi_i} \quad i = 1, \dots, k+1 \quad (\text{A.II.9})$$

#### Updating Equations

$$\frac{\partial a_\tau}{\partial \psi_i} = \frac{\partial a_{\tau/\tau-1}}{\partial \psi_i} + \frac{\partial P_{\tau/\tau-1}}{\partial \psi_i} Z_\tau f_\tau^{-1} \nu_\tau - P_{\tau/\tau-1} Z_\tau f_\tau^{-2} \frac{\partial f_\tau}{\partial \psi_i} \nu_\tau + P_{\tau/\tau-1} Z_\tau f_\tau^{-1} \frac{\partial \nu_\tau}{\partial \psi_i} \quad i = 1, \dots, k+1 \quad (\text{A.II.10})$$

$$\frac{\partial P_\tau}{\partial \psi_i} = \frac{\partial P_{\tau/\tau-1}}{\partial \psi_i} - \frac{\partial P_{\tau/\tau-1}}{\partial \psi_i} Z_\tau f_\tau^{-1} Z_\tau' P_{\tau/\tau-1} + P_{\tau/\tau-1} Z_\tau f_\tau^{-2} \frac{\partial f_\tau}{\partial \psi_i} Z_\tau' P_{\tau/\tau-1} - P_{\tau/\tau-1} Z_\tau f_\tau^{-1} Z_\tau' \frac{\partial P_{\tau/\tau-1}}{\partial \psi_i} \quad i = 1, \dots, k+1 \quad (\text{A.II.11})$$

and the initial conditions are given by:

$$\begin{aligned} \frac{\partial a_0}{\partial \psi_i} &= 0 \quad i = 1, \dots, k+1 \\ \frac{\partial P_0}{\partial \beta} &= 0 \\ \frac{\partial P_0}{\partial \rho_*} &= \begin{bmatrix} 0 & 0 \\ 0 & \frac{-2\theta_*}{(1-\rho_*^2)^2} \end{bmatrix} \\ \frac{\partial P_0}{\partial \rho_*} &= \begin{bmatrix} \frac{2(\theta_* + (\theta_* + \rho_*) \frac{\partial \theta_*}{\partial \rho_*}) (1-\rho_*^2) + 2\rho_* (1+2\rho_*\theta_* + \theta_*^2)}{(1-\rho_*^2)^2} & \frac{\partial \theta_*}{\partial \rho_*} & 0 \\ \frac{\partial \theta_*}{\partial \rho_*} & 2\theta_* \frac{\partial \theta_*}{\partial \rho_*} & 0 \\ 0 & 0 & 0 \end{bmatrix} \end{aligned}$$

### APPENDIX III

It is easy to see that

$$\frac{\partial \nu_\tau}{\partial \psi_i} = \left[ \frac{\partial \nu_\tau}{\partial \rho_*}, \frac{\partial \nu_\tau}{\partial \beta}, 0 \right] = \begin{cases} (0, -x_1^t, 0) & \tau = 1 \quad (\text{A.III.1}) \\ -(y_{\tau-1} - x_{\tau-1}^t \beta, x_\tau^t - \rho_* x_{\tau-1}^t, 0) & \tau > 1 \quad (\text{A.III.2}) \end{cases}$$

$$\frac{\partial f_\tau}{\partial \psi_i} = \left[ \frac{\partial f_\tau}{\partial \rho_*}, \frac{\partial f_\tau}{\partial \beta}, 0 \right] = \begin{cases} \left( \frac{-2\theta_*}{(1-\rho_*^2)^2}, 0, 0 \right) & \tau = 1 \quad (\text{A.III.3}) \\ 0 & \tau > 1 \quad (\text{A.III.4}) \end{cases}$$

Substituting (A.III.1-4) and (2.8-11) into (A.II.1), it follows that:

$$\begin{aligned} \frac{\partial l}{\partial \rho_*} &= \frac{-\rho_*(y_1 - x'_1\beta)^2}{\sigma_*^2} \\ &+ \frac{1}{\sigma_*^2} \sum_{r=2}^{\lfloor \frac{T}{m} \rfloor} [(y_r - x'_r\beta) - \rho_*(y_{r-1} - x'_{r-1}\beta)] (y_{r-1} - x'_{r-1}\beta) \\ &+ \frac{\rho_*}{1 - \rho_*^2} \end{aligned} \quad (\text{A.III.5})$$

$$\begin{aligned} \frac{\partial l}{\partial \beta} &= \frac{1 - \rho_*^2}{\sigma_*^2} x'_1 (y_1 - x'_1\beta) \\ &+ \frac{1}{\sigma_*^2} \sum_{r=2}^{\lfloor \frac{T}{m} \rfloor} (x_r - \rho_* x'_r)' [(y_r - x'_r\beta) - \rho_*(y_{r-1} - x'_{r-1}\beta)] \end{aligned} \quad (\text{A.III.6})$$

Substituting (2.8-11) into (A.II.2), it follows that:

$$\begin{aligned} \frac{\partial l}{\partial \sigma_*^2} &= -\frac{T}{2m\sigma_*^2} \\ &+ \frac{1 - \rho_*^2}{2\sigma_*^4} x'_1 (y_1 - x'_1\beta)^2 \\ &+ \frac{1}{\sigma_*^4} \sum_{r=2}^{\lfloor \frac{T}{m} \rfloor} [(y_r - x'_r\beta) - \rho_*(y_{r-1} - x'_{r-1}\beta)]^2 \end{aligned}$$

Substituting (2.8-11) and (A.III.1-4) into (A.II.3), it follows that:

$$\begin{aligned} I_{\rho_*\rho_*} &= \frac{1}{\sigma_*^2} \sum_{r=2}^{\lfloor \frac{T}{m} \rfloor} (y_{r-1} - x'_{r-1}\beta)^2 \\ &+ \frac{2\rho_*^2}{1 - \rho_*^2} \end{aligned} \quad (\text{A.III.7})$$

$$I_{\rho_*\beta} = \frac{1}{\sigma_*^2} \sum_{r=2}^{\lfloor \frac{T}{m} \rfloor} (x_r - \rho_* x'_r)' (y_{r-1} - x'_{r-1}\beta) \quad (\text{A.III.8})$$

$$\begin{aligned} I_{\beta\beta} &= \frac{1}{\sigma_*^2} \sum_{r=2}^{\lfloor \frac{T}{m} \rfloor} (x_r - \rho_* x'_r)(x_r - \rho_* x'_{r-1})' \\ &+ \frac{1 - \rho_*^2}{\sigma_*^2} (x_1 x'_1) \end{aligned} \quad (\text{A.III.9})$$

Substituting (A.III.1-4) and (2.8-11) into (A.II.4), it follows that:

$$I_{\rho_* \sigma_*^2} = \frac{-\rho_*}{\sigma_*^4} (y_1 - x'_1 \beta) - \frac{1}{\sigma_*^4} \sum_{r=2}^{\left[ \frac{T}{n} \right]} [(y_r - x'_r \beta) - \rho_* (y_{r-1} - x'_{r-1} \beta)] (y_{r-1} - x'_{r-1} \beta) \quad (\text{A.III.10})$$

$$I_{\beta \sigma_*^2} = \frac{1 - \rho_*^2}{\sigma_*^4} x'_1 (y_1 - x'_1 \beta) - \frac{1}{\sigma_*^4} \sum_{r=2}^{\left[ \frac{T}{n} \right]} (x_r - \rho_* x'_r)' [(y_r - x'_r \beta) - \rho_* (y_{r-1} - x'_{r-1} \beta)] \quad (\text{A.III.11})$$

Under  $H_0 : \rho = 0 \Leftrightarrow H_0; \rho_* = 0$ , it follows that:

$$\begin{aligned} \frac{\partial l}{\partial \beta}(\psi_{H_0}) &= 0 \\ \frac{\partial l}{\partial \sigma_*^2}(\psi_{H_0}) &= 0 \\ \frac{\partial l}{\partial \rho_*}(\psi_{H_0}) &= \frac{1}{\sigma_*^2} \sum_{r=2}^{\left[ \frac{T}{n} \right]} \tilde{u}_r \tilde{u}_{r-1} \end{aligned} \quad (\text{A.III.12})$$

Taking plim's in (A.III.7-11), and under appropriated regularity conditions on the  $x'(t)$ , it follows that:

$$\begin{aligned} \text{plim} \left[ \tilde{I}_{\rho_* \rho_*} - \frac{1}{\sigma_*^2} \sum_{r=1}^{\left[ \frac{T}{n} \right]} \tilde{u}_{r-1}^2 \right] &= 0 \\ \text{plim} \left[ \tilde{I}_{\beta \beta} - \frac{1}{\sigma_*^2} \sum_{r=1}^{\left[ \frac{T}{n} \right]} x_r x'_r \right] &= 0 \\ \text{plim} \tilde{I}_{\rho_* \beta} &= 0 \\ \text{plim} \tilde{I}_{\rho_* \beta} &= 0 \\ \text{plim} \tilde{I}_{\beta \sigma_*^2} &= 0 \end{aligned}$$

It follows that

$$LM = \left[ \frac{1}{\sigma_*^2} \sum_{r=1}^{\left[ \frac{T}{n} \right]} \tilde{u}_r \tilde{u}_{r-1} \right] \left[ \frac{1}{\sigma_*^2} \sum_{r=1}^{\left[ \frac{T}{n} \right]} \tilde{u}_{r-1}^2 \right]^{-1} \left[ \frac{1}{\sigma_*^2} \sum_{r=1}^{\left[ \frac{T}{n} \right]} \tilde{u}_r \tilde{u}_{r-1} \right]$$

but

$$\left[ \frac{1}{\sigma_*^2} \sum_{r=1}^{\lfloor \frac{T}{m} \rfloor} \tilde{u}_{r-1}^2 \right] = \frac{T}{m}$$

then

$$LM = \left[ \frac{T}{m} \right]^2 r_1^{*2} \left[ \frac{T}{m} \right]^{-1} = \left[ \frac{T}{m} \right] r_1^{*2}$$

where

$$r_1^{*2} = \frac{\sum_{r=1}^{\lfloor \frac{T}{m} \rfloor} \tilde{u}_r \tilde{u}_{r-1}}{\sum_{r=1}^{\lfloor \frac{T}{m} \rfloor} \tilde{u}_r^2}$$

#### APPENDIX IV

##### Proof of Proposition II:

Multiplying both sides of (A.I.2) by  $\sum_{j=1}^{m-1} L^j$ , it follows that:

$$(1 - \rho^m B)y_r = (1 - \rho^m) \left( \sum_{j=0}^{m-1} x'(t-j) \right) \beta + \omega_r \quad (\text{A.IV.1})$$

where

$$\omega_r = \left( \sum_{j=0}^{m-1} \right) \left( \sum_{j=1}^{m-1} \rho^j L^j \right) \epsilon(t) \quad (\text{A.IV.2})$$

Now (A.IV.1) can be rewritten as:

$$y_r = z_r' \beta + u_r \quad (\text{A.IV.3})$$

$$u_r = \rho_* u_{r-1} + \omega_r \quad (\text{A.IV.4})$$

where

$$z_r' = \sum_{j=0}^{m-1} x(mr-j)$$

$$\rho_* = \rho^m$$

and  $\omega_r$  is given by (A.IV.2).



Using the results of Tiao (1972) or Pereira (1986), (A.IV.4) follows an ARMA(1,1), the expression (3.3), where the AR parameter is  $\rho_*$  and the MA parameter,  $\theta_*$  and the variance of the error term,  $\sigma_*^2$ , are given by the solution of equation (3.4-6).

#### APPENDIX V

Derivating (3.9-12) with respect to  $\beta$ , it follows that:

$$\frac{\partial \nu_\tau}{\partial \beta} = \begin{cases} -z_1^t & \tau = 1 \text{ (A.V.1)} \\ -[z_\tau - \rho_* z_{\tau-1}] + \theta_* \frac{\partial f_{\tau-1}}{\partial \beta} f_{\tau-1}^{-2} \nu_{\tau-1} - \theta_* f_{\tau-1}^{-1} \frac{\partial \nu_{\tau-1}}{\partial \beta} & \tau > 1 \text{ (A.V.2)} \end{cases}$$

and

$$\frac{\partial f_\tau}{\partial \beta} = 0 \quad \text{(A.V.3)}$$

Now using (A.V.3) into (A.V.2), it follows that:

$$\frac{\partial \nu_\tau}{\partial \beta} = \begin{cases} -z_1^t & \tau = 1 \text{ (A.V.1)} \\ -[z_\tau - \rho_* z_{\tau-1}] - \theta_* f_{\tau-1}^{-1} \frac{\partial \nu_{\tau-1}}{\partial \beta} & \tau > 1 \text{ (A.V.2)} \end{cases}$$

Now derivating (3.9-12) with respect to  $\rho_*$ , it follows that:

$$\frac{\partial \nu_\tau}{\partial \rho_*} = \begin{cases} 0 & \tau = 1 \text{ (A.V.4)} \\ -[y_{\tau-1} - z_{\tau-1}^t \beta] - \frac{\partial \theta_*}{\partial \rho_*} f_{\tau-1}^{-1} \nu_{\tau-1} \\ + \theta_* \frac{\partial f_{\tau-1}}{\partial \rho_*} f_{\tau-1}^{-2} \nu_{\tau-1} - \theta_* f_{\tau-1}^{-1} \frac{\partial \nu_{\tau-1}}{\partial \rho_*} & \tau > 1 \text{ (A.V.5)} \end{cases}$$

and

$$\frac{\partial f_\tau}{\partial \rho_*} = \begin{cases} \frac{[2\theta_* + 2\rho_* \frac{\partial \theta_*}{\partial \rho_*} + 2\theta_* \frac{\partial \theta_*}{\partial \rho_*}] (1 - \rho_*^2) - 2\rho_* [1 + 2\rho_* \theta_* + \theta_*^2]}{(1 - \rho_*^2)^2} & \tau = 1 \text{ (A.V.6)} \\ 2\theta_* \frac{\partial \theta}{\partial \rho_*} (1 - f_{\tau-1}^{-1}) + \theta_*^2 \frac{\partial f_{\tau-1}}{\partial \rho_*} f_{\tau-1}^{-2} & \tau > 1 \text{ (A.V.7)} \end{cases}$$

Derivating (3.9-12) with respect to  $\sigma_*^2$ , it follows that:

$$\frac{\partial f_\tau}{\partial \sigma_*^2} = \frac{\partial \nu_\tau}{\partial \sigma_*^2} = 0$$

Then under  $H_0$ , it follows that:

$$\frac{\partial \nu_\tau}{\partial \rho_*} (\psi_{H_0}) = \begin{cases} 0 & \tau = 1 \text{ (A.V.8)} \\ -\frac{m+1}{m} [y_{\tau-1} - z_{\tau-1}^t \tilde{\beta}] & \tau > 1 \text{ (A.V.9)} \end{cases}$$

$$\frac{\partial f_{\tau}}{\partial \rho_{*}}(\psi_{H_0}) = 0 \quad \forall \tau \quad (\text{A.V.10})$$

$$\frac{\partial v_{\tau}}{\partial \beta}(\psi_{H_0}) = \begin{cases} -z_1 & \tau = 1 \quad (\text{A.V.11}) \\ -z_{\tau} & \tau > 1 \quad (\text{A.V.12}) \end{cases}$$

$$\frac{\partial f_{\tau}}{\partial \beta}(\psi_{H_0}) = 0 \quad \forall \tau \quad (\text{A.V.13})$$

Substituting (A.V.8-13) into (A.II.1), it follows that:

$$\begin{aligned} \frac{\partial l}{\partial \beta}(\psi_{H_0}) &= 0 \\ \frac{\partial l}{\partial \sigma_{*}^2}(\psi_{H_0}) &= 0 \\ \frac{\partial l}{\partial \rho_{*}}(\psi_{H_0}) &= \frac{m+1}{m\sigma_{*}^2} \sum_{\tau=2}^{\lfloor \frac{T}{m} \rfloor} \tilde{u}_{\tau} \tilde{u}_{\tau-1} \end{aligned} \quad (\text{A.V.14})$$

Taking plim's in the expressions for the information matrix and under appropriated regularity conditions on the  $x'(t)$ , it follows that:

$$\begin{aligned} \text{plim} \left[ \tilde{I}_{\rho_{*}\rho_{*}} - \left[ \frac{m+1}{m} \right]^2 \frac{1}{\sigma_{*}^2} \sum_{\tau=1}^{\lfloor \frac{T}{m} \rfloor} \tilde{u}_{\tau-1}^2 \right] &= 0 \\ \text{plim} \left[ \tilde{I}_{\beta\beta} - \frac{1}{\sigma_{*}^2} \sum_{\tau=1}^{\lfloor \frac{T}{m} \rfloor} z_{\tau} z_{\tau}' \right] &= 0 \\ \text{plim} \tilde{I}_{\rho_{*}\beta} &= 0 \\ \text{plim} \tilde{I}_{\beta\rho_{*}} &= 0 \\ \text{plim} \tilde{I}_{\beta\sigma_{*}^2} &= 0 \end{aligned}$$

It follows that

$$LM = \left[ \left[ \frac{m+1}{m} \right] \frac{1}{\sigma_{*}^2} \sum_{\tau=1}^{\lfloor \frac{T}{m} \rfloor} \tilde{u}_{\tau} \tilde{u}_{\tau-1} \right] \left[ \left[ \frac{m+1}{m} \right]^2 \frac{1}{\sigma_{*}^2} \sum_{\tau=1}^{\lfloor \frac{T}{m} \rfloor} \tilde{u}_{\tau-1}^2 \right]^{-1} \left[ \left[ \frac{m+1}{m} \right] \frac{1}{\sigma_{*}^2} \sum_{\tau=1}^{\lfloor \frac{T}{m} \rfloor} \tilde{u}_{\tau} \tilde{u}_{\tau-1} \right]$$

but

$$\left[ \frac{1}{\sigma_{*}^2} \sum_{\tau=1}^{\lfloor \frac{T}{m} \rfloor} \tilde{u}_{\tau-1}^2 \right] = \frac{T}{m}$$

then

$$LM = \left[ \frac{T}{m} \right]^2 r_1^{*2} \left[ \frac{T}{m} \right]^{-1} = \left[ \frac{T}{m} \right] r_1^{*2}$$

where

$$r_1^{*2} = \frac{\sum_{r=1}^{\left[ \frac{T}{m} \right]} \tilde{u}_r \tilde{u}_{r-1}}{\sum_{r=1}^{\left[ \frac{T}{m} \right]} \tilde{u}_r^2}$$

and  $\tilde{u}_r = y_r - z_r' \tilde{\beta}$  and  $\tilde{\beta}$  is given by (3.3):

## REFERENCES

- ANSLEY, C. I. & KOHN, R. (1985) *Estimation, Filtering and Smoothing in State Space Models With Incompletely Specified Initial Conditions*, Annals of Statistics, ( to appear).
- BREUSCH, T. S. (1978) *Testing for Autocorrelation in Dynamic Linear Models*, Australian Economic Paper, **17**, 334 - 355.
- BREUSCH, T. S. & PAGAN, A. R. (1980) *The Lagrange Multiplier Test and Its Applications to Model Specification in Econometrics*, Review of Economic Studies, **47**, 239 - 254.
- DUDOUR, J - M & DAGENAIS, M. G. (1985) *Durbin - Watson Tests for Serial Correlation in Regressions With Missing Observations*, Journal of Econometrics, **27**, 371 - 381.
- DURBIN, J. & WATSON, G. S. (1950) *Testing for Serial Correlation in Least Squares Regressions I*, Biometrika, **37**, 409 - 426.
- DURBIN, J. & WATSON, G. S. (1951) *Testing for Serial Correlation in Least Squares Regressions II*, Biometrika, **38**, 159 - 178.
- DURBIN, J. & WATSON, G. S. (1971) *Testing for Serial Correlation in Least Squares Regressions III*, Biometrika, **58**, 1 - 19.
- DURBIN, J. (1970) *Testing for Serial Correlation in Least Squares Regressions When Some of the Regressors are Lagged Dependent Variables*, Econometrica, **38**, 410 - 421.
- ENGLE, R. & WATSON, M. (1981) *A One Factor Multivariate Time Series Model of Metropolitan Wage Rates*, Journal of the American Statistical Association, **76**, 774 - 781.
- GODFREY, L. G. (1978) *Testing Against general Autoregressive and Moving Average Error Models When the Regressors Include Lagged Dependent variables*. Econometrica, **46**, 1293 - 1301.
- HARVEY, A. C. (1981) Time Series Models. Deddington: Philip Allan.
- PAGAN, A. (1978) *A Unified Approach to Estimation and Inference for Stochastically Varying Coefficient Regression Models*, Core Discussion Paper n° 7814.
- PEREIRA, P. L. V. (1985) *Testing for Serial Correlation in Regression Models with Missing Observations*. Paper presented at the I Econometrics and Time Series Workshop, July 1985, IMPA, Rio de Janeiro, Brasil.
- PEREIRA, P. L. V. (1986) *The Loss of Efficiency in Estimation of Stationary Economic Time Series Models With Missing Observations*, submitted to Journal of Time Series Analysis.
- ROBINSON, P. M. (1985) *Testing for Serial Correlation in Regression with Missing Observations*, Journal of the Royal Statistical Society, Series B, **47**, 429 - 437.
- SAVIN, N. E. & WHITE, K. J. (1978) *Testing for Serial Correlation With Missing Observations*, Econometrica, **46**, 59 - 67.
- TIAO, G. C. (1972) *Asymptotic Behaviour of Time Series Aggregated*, Biometrika, **59**, 523 - 531.

Textos para Discussão Interna editados a partir de janeiro de 84

- Nº 62 - "A Crise do Setor Externo e o Ajustamento requerido pelas Opções de Política Econômica", Milton Pereira de Assis, Janeiro 1984, 37 p.
- Nº 63 - "O Papel Atual da Fronteira Agrícola", Maria Beatriz de Albuquerque David, Fevereiro 1984, 19 p.
- Nº 64 - "As Negociações Financeiras Internacionais do Brasil Pós-FMI", José Cláudio Ferreira da Silva e Maria Helena T. T. Horta, Fevereiro 1984, 34 p.
- Nº 65 - "Insumos Modernos na Agricultura Brasileira", Cláudio Roberto Contador e Léo da Rocha Ferreira, Janeiro 1984, 159 p.
- Nº 66 - "Política Tarifária das Empresas de Saneamento: Uma Avaliação da Progressividade dos Preços", Thompson Almeida Andrade, Fevereiro 1984, 24 p.
- Nº 67 - "A Economia Brasileira: Uma Interpretação Econométrica", Versão IV, Cláudio Roberto Contador, Março 1984, 99 p.
- Nº 68 - "Comentários sobre o livro de William R. Cline, "International Debt and Stability of the World Economy", Marcelo de Moura Lara Resende, Maio 1984, 23 p.
- Nº 69 - "Crescimento Industrial, Ajuste Estrutural e Exportações de Manufaturados: Notas para a Definição de Uma Estratégia de Longo Prazo para a Economia Brasileira", Regis Bonelli e José Cláudio Ferreira da Silva, Novembro 1984, 38 p.
- Nº 70 - "Projeções da População Total, Urbano-Rural e Economicamente Ativa segundo Algumas Alternativas de Crescimento Demográfico", Maria Helena F. T. Henriques, Janeiro 1985, 56 p.

## II

- Nº 71 - "Crescimento Econômico e Oferta de Alimentos no Brasil", Gervásio Castro de Rezende, Janeiro 1985, 39 p.
- Nº 72 - "A Política Agrícola e a Diminuição do Subsídio do Crédito Rural", Gervásio Castro de Rezende, Janeiro 1985, 23 p.
- Nº 73 - "Tendências a Médio Prazo da Previdência Social Brasileira: Um Modelo de Simulação", Francisco E.B. de Oliveira, Kaizô Iwakami Beltrão, Maria Helena F.T. Henriques, Afonso Sant'Anna Bevilaqua, Alexandre Goretkin Neto, Janeiro 1985, 299 p.
- Nº 74 - "Balanço de Pagamentos Brasileiro: Um Modelo de Simulação", Ajax Reynaldo Bello Moreira, Janeiro 1985, 77 p.
- Nº 75 - "Interação entre Mercados de Trabalho e Razão entre Salários Rurais e Urbanos no Brasil", Gervásio Castro de Rezende, Março 1985, 35 p.
- Nº 76 - "Considerações sobre uma Possível Reforma Tributária no Brasil", Cláudia Cunha Campos Eris, Março 1985, 29 p.
- Nº 77 - "Migrações Internas e Pequena Produção Agrícola na Amazônia: Uma Análise da Política de Colonização do INCRA", Anna Luiza Ozorio de Almeida, Maio 1985, 97 p.
- Nº 78 - "Estrutura Industrial e Exportação de Manufaturados: Brasil, 1978", Helson C. Braga e Edson P. Guimarães, Julho 1985, 29 p.
- Nº 79 - "A Restrição Externa à Retomada do Crescimento: Avaliação e Recomendações de Política", Helson C. Braga, Setembro 1985, 42 p.
- Nº 80 - "Foreign Direct Investment in Brazil: Its Role, Regulation and Performance", Helson C. Braga, Outubro 1985, 41 p.

### III

- Nº 81 - "Déficit de "Caixa" do Governo Federal: Metodologia e Resultados em 1985", Carlos von Doellinger, Novembro 1985, 16 p.
- Nº 82 - "Déficit e Dívida: Tendências e Implicações", Carlos von Doellinger, Novembro 1985, 12 p.
- Nº 83 - "As Interligações Setoriais na Economia Brasileira em 1975", José W. Rossi, Maristela Sant'Anna e Samuel Sidsamer, Novembro 1985, 30 p.
- Nº 84 - "Mensuração da Eficiência Produtiva na Indústria Brasileira: 1980", Helson C. Braga e José W. Rossi, Novembro 1985, 34 p.
- Nº 85 - "Fundos Sociais", Fernando A. Rezende da Silva e Beatriz Azeredo da Silva, Janeiro 1986, 29 p.
- Nº 86 - "Optimal Foreign Borrowing in a Multisector Dynamic Equilibrium Model: A Case Study for Brazil", Octávio A. F. Tourinho, Janeiro 1986, 47 p.
- Nº 87 - "Proposta de Diretrizes Preliminares para Uma Política de Abastecimento", Maria Beatriz de A. David, Março 1986, 44 p.
- Nº 88 - "Os Impactos da Política de Comercialização Agrícola sobre a Produção e os Preços. Uma Análise da Literatura e Algumas Evidências Empíricas", Maria Beatriz de A. David e Luis Alberto de L.C. Ribeiro, Março 1986, 49 p.
- Nº 89 - "Distribuição de Renda: 1970/1980", José W. Rossi, Maio 1986, 17 p.
- Nº 90 - "Balança Comercial e Dinâmica da Desvalorização Cambial no Brasil, 1970/84", Helson C. Braga e José W. Rossi, Maio 1986, 20 p.

IV

- Nº 91 - "Algumas Considerações sobre os Efeitos da Reforma Monetária no Campo Social: Seguro-Desemprego e Previdência Social", Francisco E.B. de Oliveira, Kaizô Iwakami Beltrão e Marco Aurélio de Sã Ribeiro (estagiário), Maio 1986, 16 p.
- Nº 92 - "Modelos de Previsão para Séries de Produção e Preços: Metodologia Bayesiana e Box-Jenkins para Séries Temporais", Gutemberg H. Brasil, Hélio S. Migon, Reinaldo C.Souza, Sérgio S. Portugal, Maio 1986, 63 p.
- Nº 93 - "O Controle de Preços dos Alimentos e seus Efeitos sobre a Produção e o Abastecimento. Algumas Considerações para o Ano de 1986", Maria Beatriz de A. David, Junho 1986, 39 p.
- Nº 94 - "Previsão da Inflação e Produção Industrial Pós-Choque via Análise de Intervenção", H.S. Migon e G.H. Brasil, Julho 1986, 18 p.
- Nº 95 - "Exacerbação do Consumo e Salário Médio:Evidências sobre o Efeito-Sincronização", Ricardo Cicchelli Velloso, setembro 1986, 20 p.
- Nº 96 - "The Demand for Money in Brazil Revisited", José Rossi, Outubro 1986, 24 p.
- Nº 97 - "O Programa de Estabilização Econômica e o Poder de Compra do Salário Mínimo", Daniel A.Ribeiro de Oliveira e Ricardo Cicchelli Velloso, outubro 1986, 19 p.
- Nº 98 - "Formação de Expectativas num Contexto de Inflação Baixa e Alta Incerteza", Fabio Giambiagi, Outubro 1986, 38 p.



- Nº 99 - "Progresso Técnico na Indústria Brasileira: Indicadores e Análise de seus Fatores Determinantes", Helson C.Braga e Virene Matesco, Outubro 1986, 71 p.
- Nº 100 - "As Migrações Internas e a Previdência Social", Maria Helena F.T.Henriques e Kaizô I. Beltrão, Outubro 1986, 59 p.

O INPES edita ainda as seguintes publicações: Pesquisa e Planejamento Econômico (quadrimestral), desde 1971; Literatura Econômica (bimestral), desde 1977; Coleção Relatórios de Pesquisa; Série Textos para Discussão do Grupo de Energia (TDE); Série Monográfica; Série PNPE e Série Estudos de Política Industrial e Comércio Exterior (EPICO).